

Self-trapping phenomenon in (2+1)-dimensional quantum electrodynamics and its application to high- T_c cuprates with Néel ordering

T. Morinari¹

¹*Yukawa Institute for Theoretical Physics, Kyoto University Kyoto 606-8502, Japan*
(Dated: February 2, 2008)

The strong coupling effect in the (2+1)-dimensional quantum electrodynamics (QED₃) description of the $S = 1/2$ Heisenberg antiferromagnet is studied in terms of a canonical transformation which has been used in the small polaron theory in electron-phonon systems. In the Néel ordered phase, we show that the Dirac fermions are localized and the spectral function is of the Gaussian form due to the coupling to the longitudinal gauge field. The width of the broad line shape is $\sim 3J$ with J the superexchange interaction energy and the localization length is $\sim 2a$, with a the lattice constant.

PACS numbers:

I. INTRODUCTION

One of the most fundamental question in the physics of high- T_c cuprates is about the description of the doped holes. The simplest situation would be realized in the single hole doped system. Experimentally, angle resolved photoemission spectroscopy (ARPES) in the undoped compounds provides us valuable information about the nature of the excitations. The excitation spectrum was obtained by Wells *et al.*^{1,2,3} It was found that the band width is $\sim 2.2J \simeq 270\text{meV}$ where J is the superexchange interaction between the copper site spins. This band width is much smaller than the tight-binding value of $8t \simeq 2.8\text{eV}$ which is determined by the transfer integral parameter, t . Furthermore, the observed spectra exhibit a quite broad feature. In Ref.⁴, it is shown that the spectra do not have a conventional Lorentzian form. The spectra rather have a Gaussian form. These observations suggest that the quasiparticle excitations observed in the undoped cuprates are quite different from conventional Fermi liquid quasiparticles.

In order to understand ARPES experiments in the undoped compounds, first we need to figure out the elementary excitation whose excitation spectrum obeys the dispersion observed experimentally. The next step is to consider a mechanism of spectral shape broadening that would arise from coupling to a boson mode. Concerning the first point, it was shown that the slave-fermion approach to the t-t'-t''-J model reproduces the dispersion.^{5,6} (It is known that the simple t-J model fails to reproduce the dispersion along $(\pi, 0)$ to $(0, \pi)$.^{7,8}) However, it has been shown⁵ that the coupling to antiferromagnetic spin-wave modes does not lead to a broad line shape. Recently Mischenko and Nagaosa⁹ considered a coupling to an optical phonon. They numerically summed over Feynman diagrams including vertex corrections for phonons. It was argued that the quasiparticle in the t-J model is in the strong coupling regime that leads to a broad line shape.

In this paper, we consider another possibility. We consider the quasiparticles in the π -flux phase¹⁰ and the effect of coupling to U(1) gauge field fluctuations. A similar problem within the slave-fermion approach to the t-J model was discussed by Auerbach and Larson.¹¹ It was argued that the single hole forms a small polaron due to the coupling to the gauge field. Here we consider a similar problem based on the π -flux phase with dynamically induced mass. The π -flux phase was proposed by Affleck and Marston¹⁰ from a mean field theory of the $S = 1/2$ antiferromagnetic Heisenberg model based on a fermionic representation of the spin $S = 1/2$. Fluctuations about the state can be described by a U(1) gauge field. The dispersion of the quasiparticle in the π -flux phase is in good agreement with the experimentally obtained dispersion as argued by Laughlin.¹² The effective theory is described by quantum electrodynamics in (2+1)-dimension, which is called QED₃.^{13,14}

Although the π -flux phase is different from the Néel ordering phase, believed to describe the Néel ordering state^{13,14} is the state that includes the dynamically induced mass term due to the coupling to the gauge field. The excitation spectra near $(\pm\pi/2, \pm\pi/2)$ are given by a massive Dirac fermion. Experimentally the Dirac fermion mass is estimated to be $\sim 1.3J$ ¹⁵ by fitting the experimentally obtained dispersion with the quasiparticle dispersion in the π -flux phase with the mass term.

Meanwhile Franz and Tešanović proposed¹⁶ a QED₃ theory starting from the d-wave superconducting state.¹⁷ The dynamical mass generation, or the spontaneous chiral symmetry breaking, is also associated with antiferromagnetic order.¹⁸ Aitchison and Mavromatos¹⁹ argued that the gauge-fermion interaction in QED₃ leads to non-Fermi liquid behavior. Rantner and Wen^{20,21} discussed changes of line shapes between the normal state and the superconducting state based on a flux phase obtained in the SU(2) slave-boson approach to the t-J model²² with QED₃ for the description of the spinons.

In the Néel ordered phase, there is strong logarithmic potential between the Dirac fermions. In fact, there is no low-lying spin 1/2 excitations^{23,24} because of the confinement associated with this strong interaction.²⁵ Therefore, in the Néel phase we need to perform a non-perturbative analysis.

In this paper, we apply to the QED₃ a canonical transformation, which has been extensively studied in the electron-phonon systems,²⁶ for the effect of the longitudinal gauge field interaction on the massive Dirac fermion. We show that the coupling to the longitudinal gauge field leads to the localization of the Dirac fermions and the spectral function is approximately given by a Gaussian form. The main assumption here is that a doped hole excite a quasiparticle in the π -flux state. One way to interpret this is to consider the slave-boson approach to the t-J model. Another way is proposed in Ref.²⁷ where a half-skyrmion spin texture is created by doped holes. The excitation spectrum of the half-skyrmion spin texture is the same as that of the π -flux phase. The mass term of the quasiparticles in the π -flux state is the excitation gap of the half-skyrmion spin texture. The effective action of the half-skyrmion spin textures is shown to be described by QED₃ by applying a duality mapping.²⁷

The rest of the paper is organized as follows. In Sec.II, we describe the QED₃ action and the application of the canonical transformation. In Sec.III, we calculate the retarded Green's function of the massive Dirac fermion. Section IV is devoted to the conclusion.

II. MODEL

In the continuum, the QED₃ action for the $S = 1/2$ quantum Heisenberg antiferromagnet^{13,14} is written as

$$\begin{aligned} S &= \int d^3x \bar{\psi}(x) (i\gamma^\mu D_\mu - mc_{sw}\sigma_3) \psi(x) - \frac{1}{4e_A^2} \int d^3x F_{\mu\nu} F^{\mu\nu} \\ &= S_0 + S_{\text{int}} + S_A, \end{aligned} \quad (1)$$

where $D_\mu = \partial_\mu + iA_\mu$ is the covariant derivative. (The derivation of the action in the π -flux phase is given in appendix A.) The Dirac fermion fields describe the quasiparticles in the spin system. In the absence of the doped holes, the ground state is that all the negative energy states of the Dirac fermions are occupied. While the positive energy states are empty. Upon hole doping, hole states are created in the band of the negative energy states. In writing down the action, we have assumed that there is the Néel ordering. In the presence of the Néel ordering, the mass m is nonzero.^{13,14} The dynamics of the gauge field arises from the fermion polarization.¹⁴ Due to the mass term the gauge field action takes the form of the Maxwell action. The gauge charge e_A is related to m through $e_A^2 = 3\pi mc_{sw}$.¹⁴

For static particles, the scalar potential $\phi \equiv A_0/e_A$ is calculated by the following equation:

$$\nabla^2 \phi(\mathbf{r}) = -e_A \rho(\mathbf{r}), \quad (2)$$

where $\rho(\mathbf{r})$ represents the charge density. The potential between a particle and hole is given by

$$V(r) = -\frac{e_A^2}{2\pi} \ln \frac{r}{r_0}. \quad (3)$$

Here r_0 is the short distance cutoff scale which is on the order of the lattice constant. Note that the potential $V(r)$ has a form of a logarithmically confining potential.

We consider the effect of the scalar potential on a dynamical hole. In solid state physics, the Coulomb gauge is often used for the analysis of the electromagnetic gauge field. In the Coulomb gauge, the interaction mediated by the scalar potential is instantaneous in time. While in the Lorentz gauge, the retardation effect in the interaction is explicit. For the electromagnetic gauge field, the speed of light c is much faster than vectors of excitations in the sample. Therefore, the retardation effect is negligible. However, for the QED₃ theory of (1), the speed of the gauge field propagation is c_{sw} that is much smaller than c . The retardation effect can play an important role. To include this effect, we take the Lorentz gauge. In the Lorentz gauge, the action of the gauge field is

$$S_A = \frac{c_{sw}}{2e_A^2} \int dt \int d^2\mathbf{r} \sum_{\mu=0,1,2} \left[\frac{1}{c_{sw}^2} (\partial_t A_\mu)^2 - (\partial_x A_\mu)^2 - (\partial_y A_\mu)^2 \right]. \quad (4)$$

The vector potential obeys the following equation,

$$\frac{1}{c_{sw}^2} \partial_t^2 \mathbf{A} - \partial_x^2 \mathbf{A} - \partial_y^2 \mathbf{A} = -\mathbf{j}. \quad (5)$$

Note that the equation for the longitudinal component of the vector potential is derived from the equation for the scalar potential in the Lorentz gauge. Therefore, the independent degrees of freedom are the transverse component of the vector potential and the scalar potential. The Hamiltonian for the gauge field is

$$H_A = \frac{1}{2} \sum_{q, \mu=0, t} \omega_q (a_{q\mu}^\dagger a_{q\mu} + a_{q\mu} a_{q\mu}^\dagger), \quad (6)$$

where $\omega_q = c_{sw}q$ in the continuum and we have introduced creation and annihilation operators through $A_{q\mu} = \sqrt{e_A^2 c_{sw}/(2\omega_q)} (a_{-q, \mu}^\dagger + a_{q\mu})$. Here $\mu = t$ denotes the transverse component of the vector potential.

On the square lattice, the action $S_0 + S_{\text{int}}$ has the following form²⁸

$$\begin{aligned} S_0 + S_{\text{int}} = & \int dt a^2 \sum_j \{ \bar{\psi}_j(t) i\gamma^0 \partial_t \psi_j(t) - \bar{\psi}_j(t) c_{sw} \gamma^0 A_0(j, t) \psi_j(t) - m c_{sw}^2 \bar{\psi}_j(t) \sigma_3 \psi_j(t) \\ & - \frac{c_{sw}}{2a} \bar{\psi}_j(t) i\gamma^1 \left[e^{iaA_1(j+a\hat{e}_1/2, t)} \psi_{j+a\hat{e}_1}(t) - e^{-iaA_1(j-a\hat{e}_1/2, t)} \psi_{j-a\hat{e}_1}(t) \right] \\ & - \frac{c_{sw}}{2a} \bar{\psi}_j(t) i\gamma^2 \left[e^{iaA_2(j+a\hat{e}_2/2, t)} \psi_{j+a\hat{e}_2}(t) - e^{-iaA_2(j-a\hat{e}_2/2, t)} \psi_{j-a\hat{e}_2}(t) \right] \}. \end{aligned} \quad (7)$$

The above action contains the gauge field interactions arising from both the transverse part and the longitudinal part of the gauge field. The transverse component of the vector potential is important for the calculation of the quasiparticle lifetime. Because the scattering process associated with the transverse component is due to emissions or absorptions of the propagating gauge fields, or “photons.” As demonstrated in Refs.^{29,30}, the coupling to the transverse vector potential leads to a T -linear resistivity law. However, the interaction of the holes with the transverse vector potential does not creat a trapping potential, and is described by a self-energy effect. Therefore, the spectrum has the form of Lorentzian which is inconsistent with the ARPES experiments in the undoped compounds. Here we focus on the self-trapping phenomenon associated with the scalar potential, and neglect the transverse component of the vector potential:

$$\begin{aligned} H_{\text{int}} &= c_{sw} a^2 \sum_j \bar{\psi}_j \gamma^0 A_0(j) \psi_j \\ &= \frac{c_{sw} a^2}{\sqrt{\Omega}} \sum_j \sum_{\mathbf{q}} \psi_j^\dagger \psi_j e^{i\mathbf{q} \cdot \mathbf{R}_j} A_{\mathbf{q}}. \end{aligned} \quad (8)$$

The interaction H_{int} contains the terms that contribute to the infrared (IR) behavior of QED₃^{21,31,32} and the other terms. As is well-known in QED₃, the dynamical mass generation is associated with the IR behavior.³³ For the analysis, we need a non-perturbative scheme, such as a $1/N$ expansion with N the number of Dirac fermion species. For discussion of the dynamical mass generation, the Schwinger-Dyson equation has been studied.³² In the analysis of the Schwinger-Dyson equation, it is necessary to figure out the form of the vertex part. For the IR region, one needs to properly choose the form of the vertex part for the discussion of the dynamical mass generation. While for the other region, the bare vertex form is known to be sufficient for the analysis of the system.³⁴ Here we are interested in the interaction with the momentum which is not in the IR regime. The IR behavior of the system is understood to be associated with the mass generation. Therefore, we introduce an IR cut off q_0 , which is much smaller than the ultraviolet cut off $\Lambda \sim \pi/a$, in the following calculations.

We consider a canonical transformation:²⁶

$$\bar{H} = \exp(s) H \exp(-s), \quad (9)$$

where

$$s = \sum_q \frac{M_q}{\omega_q} (a_{-q}^\dagger - a_q), \quad (10)$$

with $a_q^\dagger = a_{q0}^\dagger$ and $a_q = a_{q0}$, and

$$M_q = \frac{a^2}{\sqrt{\Omega}} \sum_j \sqrt{\frac{e_A^2 c_{sw}^3}{2\omega_q}} \psi_j^\dagger \psi_j \exp(i\mathbf{q} \cdot \mathbf{R}_j). \quad (11)$$

The bare vertex is used here as discussed above. Hereafter the summation in the momentum space concerning the gauge fields is to be performed for $q_0 < |\mathbf{q}| < \Lambda$.

Using the following result,

$$[s, \psi_j] = -\frac{a^2 e_A}{\sqrt{2\Omega}} \sum_{\mathbf{q}} \left(\frac{c_{\text{sw}}}{\omega_q} \right)^{3/2} e^{i\mathbf{q} \cdot \mathbf{R}_j} (a_{-q}^\dagger - a_{\mathbf{q}}) \psi_j, \quad (12)$$

we find that the fermion field ψ_j is transformed into

$$e^s \psi_j e^{-s} = X_j \psi_j, \quad (13)$$

where

$$X_j = \exp \left[-\frac{a^2 e_A}{\sqrt{2\Omega}} \sum_{\mathbf{q}} \left(\frac{c_{\text{sw}}}{\omega_q} \right)^{3/2} \exp(i\mathbf{q} \cdot \mathbf{R}_j) (a_{-q}^\dagger - a_{\mathbf{q}}) \right]. \quad (14)$$

The transformed Hamiltonian reads

$$\overline{H} = \overline{H}_0 + \overline{H}_{\text{int}}, \quad (15)$$

$$\begin{aligned} \overline{H}_0 = & -\frac{c_{\text{sw}} a}{2} \sum_{j, \mu=1,2} \overline{\psi}_j \gamma^\mu (\psi_{j+a\hat{e}_\mu} - \psi_{j-a\hat{e}_\mu}) - m c_{\text{sw}}^2 \sum_j \overline{\psi}_j \sigma_3 \psi_j \\ & + \frac{1}{2} \sum_{\mathbf{q}} \omega_q (a_{\mathbf{q}}^\dagger a_{\mathbf{q}} + a_{\mathbf{q}} a_{\mathbf{q}}^\dagger), \end{aligned} \quad (16)$$

$$\overline{H}_{\text{int}} = - \sum_{\mathbf{q}} \frac{M_{-\mathbf{q}} M_{\mathbf{q}}}{\omega_q}, \quad (17)$$

where we have omitted the band narrowing factor $X_j^\dagger X_{j \pm a\hat{e}_\mu}$, in the hopping term. (The estimation is similar to that of eqs.(38) and (39) below. See Appendix B.) $\overline{H}_{\text{int}}$ describes the logarithmic interaction between the Dirac fermions. In the following, we consider the effect of the longitudinal gauge field on the Dirac fermion spectrum given by $\overline{H}_0$.

III. CALCULATION OF THE RETARDED GREEN'S FUNCTION

The retarded Green's function may be given by

$$G_R(i, j; t) = -i\theta(t) \langle \psi_i(t) \overline{\psi}_j(0) + \overline{\psi}_j(0) \psi_i(t) \rangle \quad (18)$$

We separate this quantity into two terms,

$$G_R(i, j; t) = G_1(i, j; t) + G_2(i, j; t), \quad (19)$$

where

$$G_1(i, j; t) = -i\theta(t) \langle \psi_i(t) \overline{\psi}_j(0) \rangle, \quad (20)$$

$$G_2(i, j; t) = -i\theta(t) \langle \overline{\psi}_j(0) \psi_i(t) \rangle. \quad (21)$$

In the following calculations, we only consider the Dirac fermions with the spin up component and residing near $(\pi/2, \pi/2)$ in the momentum space. We redefine the Dirac fermion field $\psi(x)$ so that it represents the spin up Dirac fermion and describes the components near $(\pi/2, \pi/2)$. For the γ matrices, we may use the reduced form of γ matrices: $\gamma^0 \rightarrow \tau_3$, $\gamma^1 \rightarrow i\tau_2$, and $\gamma^2 \rightarrow i\tau_1$ with τ_μ the Pauli matrices because we only consider the two components of the

Dirac fermion fields. (The calculations below are equally applied to the Dirac fermions with the spin down component and/or residing near $(-\pi/2, \pi/2)$.) For $t > 0$, by performing the canonical transformation we have

$$\begin{aligned}
iG_1(i, j; t) &= \langle \psi_i(t) \bar{\psi}_j(0) \rangle \\
&= \frac{\text{Tr} e^{-\beta \bar{H}_0} e^{i \bar{H}_0 t} X_i \psi_i e^{-i \bar{H}_0 t} \bar{\psi}_j X_j^\dagger}{\text{Tr} e^{-\beta \bar{H}_0}} \\
&= \left\langle \exp \left[-\frac{e_A}{\sqrt{2\Omega}} \sum_q \frac{1}{\omega_q^{3/2}} \left(a_{-q}^\dagger e^{i\omega_q t} - a_q e^{-i\omega_q t} \right) e^{iq \cdot R_i} \right] \right. \\
&\quad \times \frac{1}{\Omega} \sum_{k, k'} e^{-i\gamma_0(i \sin k_1 \gamma_1 + i \sin k_2 \gamma_2 + m)t} \psi_k \bar{\psi}_{k'} e^{ik \cdot R_i} e^{-ik' \cdot R_j} \\
&\quad \left. \times \exp \left[\frac{e_A}{\sqrt{2\Omega}} \sum_q \frac{1}{\omega_q^{3/2}} \left(a_{-q}^\dagger - a_q \right) e^{iq \cdot R_j} \right] \right\rangle, \tag{22}
\end{aligned}$$

where we have used

$$e^{i \bar{H}_0 t} \psi_k e^{-i \bar{H}_0 t} = \exp[i(-\tau_1 \sin k_1 + \tau_2 \sin k_2 + \tau_3 m)t] \psi_k. \tag{23}$$

(Hereafter we set $a = 1$ and $c_{\text{sw}} = 1$.) The matrix in the exponential is diagonalized by the following unitary matrix:

$$U_k = \frac{1}{\sqrt{2\varepsilon_k(\varepsilon_k + m)}} \begin{pmatrix} \varepsilon_k + m & \sin k_1 + i \sin k_2 \\ -\sin k_1 + i \sin k_2 & \varepsilon_k + m \end{pmatrix}, \tag{24}$$

$$U_k^\dagger = \frac{1}{\sqrt{2\varepsilon_k(\varepsilon_k + m)}} \begin{pmatrix} \varepsilon_k + m & -\sin k_1 - i \sin k_2 \\ \sin k_1 - i \sin k_2 & \varepsilon_k + m \end{pmatrix}, \tag{25}$$

with $\varepsilon_k = \sqrt{\sin^2 k_1 + \sin^2 k_2 + m^2}$, and then,

$$\begin{aligned}
iG_1(i, j; t) &= \left\langle \exp \left[-\frac{e_A}{\sqrt{2\Omega}} \sum_q \frac{1}{\omega_q^{3/2}} \left(a_{-q}^\dagger e^{i\omega_q t} - a_q e^{-i\omega_q t} \right) e^{iq \cdot R_i} \right] \right. \\
&\quad \times \exp \left[-\frac{e_A}{\sqrt{2\Omega}} \sum_q \frac{1}{\omega_q^{3/2}} \left(a_{-q}^\dagger - a_q \right) e^{iq \cdot R_j} \right] \Bigg\rangle \\
&\quad \times \frac{1}{\Omega} \sum_k U_k \begin{pmatrix} [1 - f(\varepsilon_k)] e^{-i\varepsilon_k t} & 0 \\ 0 & e^{i\varepsilon_k t} f(\varepsilon_k) \end{pmatrix} U_k^\dagger e^{ik \cdot (R_i - R_j)}. \tag{26}
\end{aligned}$$

Using the following formula,

$$e^{Aa^\dagger + Ba} e^{Ca^\dagger + Da} = e^{\frac{1}{2}(AB+CD)+BC} e^{(A+C)a^\dagger} e^{(B+D)a}, \tag{27}$$

$$\langle e^{Aa^\dagger} e^{Ba} \rangle = e^{AB \langle a^\dagger a \rangle}, \tag{28}$$

we obtain

$$\begin{aligned}
&\left\langle \exp \left[-\frac{e_A}{\sqrt{2\Omega}} \sum_q \frac{1}{\omega_q^{3/2}} \left(a_{-q}^\dagger - a_q \right) e^{iq \cdot R_j} \right] \exp \left[\frac{e_A}{\sqrt{2\Omega}} \sum_q \frac{1}{\omega_q^{3/2}} \left(a_{-q}^\dagger e^{i\omega_q t} - a_q e^{-i\omega_q t} \right) e^{iq \cdot R_j} \right] \right\rangle \\
&= \exp \left[-\frac{e_A^2}{2\Omega} \sum_q \frac{1}{\omega_q^3} \left(1 - e^{i\omega_q t} e^{-iq \cdot (R_i - R_j)} \right) \right] \exp \left[-\frac{e_A^2}{\Omega} \sum_q \frac{1}{\omega_q^3} (1 - \cos[\omega_q t - q \cdot (R_i - R_j)]) n(\omega_q) \right] \\
&= \exp \left[i \frac{e_A^2}{2\Omega} \sum_q \frac{1}{\omega_q^3} \sin[\omega_q t - q \cdot (R_i - R_j)] \right] \\
&\quad \times \exp \left[-\frac{e_A^2}{2\Omega} \sum_q \frac{1}{\omega_q^3} \coth \frac{\beta \omega_q}{2} (1 - \cos[\omega_q t - q \cdot (R_i - R_j)]) \right] \tag{29}
\end{aligned}$$

Since $\epsilon_k > m \simeq J$, and we are interested in the temperature region where $\beta J \gg 1$, we may assume that $f(\epsilon_k) \ll 1$. Thus,

$$\begin{aligned} iG_1(i, j; t) \simeq & \exp \left[i \frac{e_A^2}{2\Omega} \sum_q \frac{1}{\omega_q^3} \sin [\omega_q t - q \cdot (R_i - R_j)] \right] \\ & \times \exp \left[-\frac{e_A^2}{2\Omega} \sum_q \frac{1}{\omega_q^3} \coth \frac{\beta \omega_q}{2} (1 - \cos [\omega_q t - q \cdot (R_i - R_j)]) \right] \\ & \times \frac{1}{\Omega} \sum_k e^{i\varepsilon_k t} e^{ik \cdot (R_i - R_j)} \frac{1}{2\varepsilon_k} \begin{pmatrix} m + \varepsilon_k & -\sin k_1 - i \sin k_2 \\ -\sin k_1 + i \sin k_2 & \varepsilon_k - m \end{pmatrix} \end{aligned} \quad (30)$$

Performing similar calculations, we obtain

$$\begin{aligned} iG_2(i, j; t) \simeq & \exp \left[i \frac{e_A^2}{2\Omega} \sum_q \frac{1}{\omega_q^3} \sin [\omega_q t - q \cdot (R_i - R_j)] \right] \\ & \times \exp \left[-\frac{e_A^2}{\Omega} \sum_q \frac{1}{\omega_q^3} \coth \frac{\beta \omega_q}{2} (1 - \cos [\omega_q t - q \cdot (R_i - R_j)]) \right] \\ & \times \frac{1}{\Omega} \sum_k \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} e^{-i\varepsilon_k t} e^{ik \cdot (R_i - R_j)} \end{aligned} \quad (31)$$

Finally, we obtain

$$\begin{aligned} iG_R(i, j; t) \simeq & \exp \left[-\frac{e_A^2}{\Omega} \sum_q \frac{1}{\omega_q^3} \coth \frac{\beta \omega_q}{2} (1 - \cos [\omega_q t - q \cdot (R_i - R_j)]) \right] \\ & \times \frac{1}{\Omega} \sum_k \begin{pmatrix} e^{i\varepsilon_k t} e^{ik \cdot (R_i - R_j)} f(i, j; t) & 0 \\ 0 & e^{-i\varepsilon_k t} e^{ik \cdot (R_i - R_j)} f^*(i, j; t) \end{pmatrix}, \end{aligned} \quad (32)$$

with

$$f(i, j; t) = \exp \left[i \frac{e_A^2}{2\Omega} \sum_q \frac{1}{\omega_q^3} \sin (\omega_q t) \cos [q \cdot (R_i - R_j)] \right], \quad (33)$$

where we have assumed that $(m + \epsilon_k)/2\epsilon_k \simeq 1$, $(m - \epsilon_k)/2\epsilon_k \simeq 0$, and $(\sin k_1 \pm i \sin k_2)/2\epsilon_k \simeq 0$. Because for the small hole concentration, $k/m \ll 1$. Note that the factor,

$$\frac{1}{\Omega} \sum_k e^{\pm i\varepsilon_k t} e^{ik \cdot (R_i - R_j)}, \quad (34)$$

denotes the wave function of the free propagating particle.

The second exponential factor ($\equiv I$) in the last line of Eq.(29) leads to a localization effect of the Dirac fermions. I is approximated as follows,

$$I \simeq -\frac{e_A^2}{8\pi^2} \int_0^\pi dq \int_0^{2\pi} d\theta \frac{q}{\omega_q^3} \coth \frac{\beta \omega_q}{2} (1 - \cos [\omega_q t - q \cdot (R_i - R_j)]). \quad (35)$$

From numerical estimations, we see that I has the Gaussian form as shown in Figs. 1 and 2. We find that

$$I \simeq \exp (-1.8125t^2 - 0.90625R^2). \quad (36)$$

To reproduce this Gaussian form, we make a following approximation:

$$\begin{aligned} I & \simeq -\frac{e_A^2}{4} \int \frac{d^2 \mathbf{q}}{(2\pi)^2} \frac{1}{\omega_q^3} \coth \frac{\beta \omega_q}{2} [\omega_q^2 t^2 + (q \cdot (R_i - R_j))^2] \\ & \equiv -\frac{1}{4} \Delta^2 t^2 - \frac{(R_i - R_j)^2}{\ell^2}, \end{aligned} \quad (37)$$




FIG. 1: The decay factor I as the function of $R = |\mathbf{R}_{ij}|$ for several t values. The vector $\mathbf{R}_{ij}$ is assumed to be $\mathbf{R}_{ij} = (R_{ij}, 0)$.




FIG. 2: The decay factor I as the function of t for several $R_{ij} (\equiv R)$ values. The vector $\mathbf{R}_{ij}$ is assumed to be $\mathbf{R}_{ij} = (R_{ij}, 0)$ as in Fig. 1.

where

$$\Delta^2 = e_A^2 \int \frac{d^2 \mathbf{q}}{(2\pi)^2} \frac{1}{\omega_q} \coth \frac{\beta \omega_q}{2} \quad (38)$$

$$\frac{1}{\ell^2} = \frac{e_A^2}{4} \int \frac{d^2 \mathbf{q}}{(2\pi)^2} \frac{q^2}{\omega_q^3} \coth \frac{\beta \omega_q}{2} \quad (39)$$

At low temperatures, Δ is

$$\Delta^2 \simeq \frac{e_A^2}{\pi} \int_{q_0}^{\Lambda} dq \frac{q}{\omega_q} \coth \frac{\beta \omega_q}{2} = \frac{e_A^2}{\pi} \left[(\Lambda - q_0) + 2 \sum_{n=1}^{\infty} \frac{e^{-n\beta q_0} - e^{-n\beta \Lambda}}{n\beta} \right] \simeq \frac{e_A^2}{\pi} \Lambda$$

Using the values of $Z_c = 1.17$, which was estimated from quantum Monte Carlo simulations,³⁵ and $m = 1.3J$, which was estimated experimentally,¹⁵ we obtain $\Lambda \simeq \pi/a = \sqrt{2}\pi Z_c J \simeq 5J$, $e_A^2 = 3\pi m \simeq 12J$. Thus,

$$\Delta \simeq 3J. \quad (40)$$

This value is in good agreement with the value $\Delta \simeq 2.7J$, which is obtained from Eq.(36). Therefore, the above Gaussian approximation is justified. Similarly ℓ is evaluated as

$$\frac{1}{\ell^2} \simeq \frac{e_A^2}{8\pi} \int_{q_0}^{\Lambda} dq \coth \frac{\beta\omega_q}{2} \simeq \frac{e_A^2}{8\pi} \Lambda, \quad (41)$$

$$\ell \simeq 2a. \quad (42)$$

From Eq.(36), we find $\ell \simeq a$. These result suggest that the particle is localized with the localization length of $\ell = O(a)$.

The function $f(i, j; t)$ is associated with the wave function renormalization. The argument of the exponential is

$$\sim \frac{1}{2} J t \ln \left(\frac{\Lambda}{q_0} \right) \leq \ln \left(\frac{\Lambda}{q_0} \right). \quad (43)$$

(The value of t that gives the saddle point in the frequency integration may be used in the calculation of $G_R(k, \omega)$ below.) As a rough approximation, we take $f(i, j; t)$ as a constant C . Thus,

$$iG_R(i, j; t) = \exp \left[-\frac{1}{4} \Delta^2 t^2 - \frac{|R_i - R_j|^2}{\ell^2} \right] \times \begin{pmatrix} \frac{C}{\Omega} \sum_k e^{i\varepsilon_k t} e^{ik \cdot (R_i - R_j)} & 0 \\ 0 & \frac{C^*}{\Omega} \sum_k e^{-i\varepsilon_k t} e^{ik \cdot (R_i - R_j)} \end{pmatrix}. \quad (44)$$

Performing Fourier transformation, we obtain

$$iG_R(k, \omega) = \frac{8\pi^{5/2}}{\Delta} \begin{pmatrix} C \exp [-(\omega - \epsilon_k)^2 / \Delta^2] & 0 \\ 0 & C^* \exp [-(\omega + \epsilon_k)^2 / \Delta^2] \end{pmatrix} \quad (45)$$

The spectral function is

$$-\frac{1}{\pi} \text{Im} G_R(k, \omega) = Z \frac{2\pi^{1/2}}{\Delta} \begin{pmatrix} \exp [-(\omega - \epsilon_k)^2 / \Delta^2] & 0 \\ 0 & \exp [-(\omega + \epsilon_k)^2 / \Delta^2] \end{pmatrix}, \quad (46)$$

where Z is the wave function renormalization factor given by $Z = 4\pi \text{Re} C$. Thus, the spectral function has a Gaussian form. This is consistent with the experiments.⁴ The value of the width $\Delta \sim 3J$ appears reasonable compared with the experiments. The main drawback in the present calculation is that we cannot discuss anisotropy in the momentum space observed in the experiments. Because we have started from the continuum theory. In order to discuss anisotropy, we need to formulate the theory on the lattice.

IV. CONCLUSION

To conclude, we have studied the effect of the longitudinal gauge field on the quasiparticle spectrum in the π -flux phase with the dynamically induced mass. A canonical transformation is applied for the analysis of the strong coupling effect between the massive Dirac fermions and the longitudinal gauge field. We have shown that the Dirac fermion propagator has the strong decay form that leads to the localized state. The spectral function is approximately given by a Gaussian form. The width of the broad spectral line shape is $\sim 4J$ and the localization length is $\sim 3a$.

Although we have carried out the analysis in the Néel ordering phase, in the disordered phase we expect a qualitatively different behavior because there is no strong logarithmic potential that leads to confinement. It would be interesting to study the delocalization transition associated with the Néel order destruction.

Acknowledgments

I would like to thank N. Nakai and K. Shizuya for discussions. This work was supported by Grant-in-Aid for Young Scientists(B) (17740253) and the 21st Century COE "Center for Diversity and Universality in Physics" from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan.

APPENDIX A: DERIVATION OF QED₃ ACTION IN THE π -FLUX PHASE

In this appendix we derive the QED₃ action (1) in the π -flux phase¹⁰ with the dynamically induced mass m . The $S = 1/2$ antiferromagnetic Heisenberg model is given by

$$H = J \sum_{\langle i,j \rangle} \mathbf{S}_i \cdot \mathbf{S}_j, \quad (\text{A1})$$

where the summation is taken over the nearest neighbor sites. The spin 1/2 can be represented by fermion fields,

$$\mathbf{S}_j = \frac{1}{2} \begin{pmatrix} f_{j\uparrow}^\dagger & f_{j\downarrow}^\dagger \end{pmatrix} \boldsymbol{\sigma} \begin{pmatrix} f_{j\uparrow} \\ f_{j\downarrow} \end{pmatrix}, \quad (\text{A2})$$

where $f_{j\sigma}$ and $f_{j\sigma}^\dagger$ are fermion annihilation and creation operators at site j with spin σ , respectively. The fermion operators satisfy the constraint, $\sum_\sigma f_{j\sigma}^\dagger f_{j\sigma} = 1$. In this fermion representation, the Hamiltonian reads

$$H = -\frac{J}{2} \sum_{\langle i,j \rangle} \sum_{\alpha,\beta} f_{i\alpha}^\dagger f_{j\alpha} f_{j\beta}^\dagger f_{i\beta} + i \sum_j \lambda_j \left(\sum_\alpha f_{j\alpha}^\dagger f_{j\alpha} - 1 \right) + \text{const.} \quad (\text{A3})$$

Here λ_j is a Lagrange multiplier to impose the constraint. Following Affleck and Marston,¹⁰ we take the mean fields,

$$\chi_{ij} = \sum_\alpha \langle f_{j\alpha}^\dagger f_{i\alpha} \rangle, \quad \lambda_j = \lambda. \quad (\text{A4})$$

We set $\chi_1 = \chi_{j,j+\hat{e}_x}$, $\chi_2 = \chi_{j+\hat{e}_x,j+\hat{e}_y}$, $\chi_3 = \chi_{j+\hat{e}_x+\hat{e}_y,j+\hat{e}_y}$, and $\chi_4 = \chi_{j+\hat{e}_y,j}$. Now the Hamiltonian is

$$\begin{aligned} H = & -\frac{J}{2} \sum_{j \in \text{even}} \sum_\sigma \left(\chi_1 f_{j+\hat{x},\sigma}^\dagger f_{j\sigma} + \chi_4^* f_{j+\hat{y},\sigma}^\dagger f_{j\sigma} + h.c. \right) - \frac{J}{2} \sum_{j \in \text{odd}} \left(\chi_3^* f_{j+\hat{x},\sigma}^\dagger f_{j\sigma} + \chi_2 f_{j+\hat{y},\sigma}^\dagger f_{j\sigma} + h.c. \right) \\ & + \frac{NJ}{4} (|\chi_1|^2 + |\chi_2|^2 + |\chi_3|^2 + |\chi_4|^2) + i\lambda \sum_j \left(\sum_\sigma f_{j\sigma}^\dagger f_{j\sigma} - 1 \right). \end{aligned} \quad (\text{A5})$$

Performing the Fourier transformation

$$f_{j\sigma} = \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{R}_j} f_{\mathbf{k}\sigma} = \frac{1}{\sqrt{N}} \sum'_{\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{R}_j} [f_{\mathbf{k}\sigma} + (-1)^j f_{\mathbf{k}+\mathbf{Q}\sigma}], \quad (\text{A6})$$

with $\sum'_{\mathbf{k}}$ being the summation over the half of the Brillouin zone, we obtain

$$H = -\frac{J}{2} \sum'_{\mathbf{k}} \begin{pmatrix} f_{e\mathbf{k}}^\dagger & f_{o\mathbf{k}}^\dagger \end{pmatrix} \begin{pmatrix} 0 & \kappa_{\mathbf{k}}^* \\ \kappa_{\mathbf{k}} & 0 \end{pmatrix} \begin{pmatrix} f_{e\mathbf{k}} \\ f_{o\mathbf{k}} \end{pmatrix} + \text{const.}, \quad (\text{A7})$$

where the two-component spinors $f_{e\mathbf{k}}$ and $f_{o\mathbf{k}}$ are $f_{e\mathbf{k}} = (f_{\mathbf{k}} + f_{\mathbf{k}+\mathbf{Q}})/\sqrt{2}$ and $f_{o\mathbf{k}} = (f_{\mathbf{k}} - f_{\mathbf{k}+\mathbf{Q}})/\sqrt{2}$, and $\kappa_{\mathbf{k}} = \chi_1 e^{-ik_x} + \chi_3 e^{ik_x} + \chi_2^* e^{ik_y} + \chi_4^* e^{-ik_x}$.

In the Néel ordering phase, there is the staggered magnetization term

$$\sum_j m_{\text{st}} (-1)^j f_j^\dagger \sigma_3 f_j = m_{\text{st}} \sum'_{\mathbf{k}} \left(f_{e\mathbf{k}}^\dagger \sigma_3 f_{e\mathbf{k}} - f_{o\mathbf{k}}^\dagger \sigma_3 f_{o\mathbf{k}} \right), \quad (\text{A8})$$

where σ_3 is a Pauli matrix in spin space. Following Affleck and Marston,¹⁰ we take $\chi_1 = \chi_3 = |\chi|$ and $\chi_2 = \chi_4 = i|\chi|$. Then,

$$H = \sum'_{\mathbf{k}} \begin{pmatrix} f_{e\mathbf{k}}^\dagger & f_{o\mathbf{k}}^\dagger \end{pmatrix} \begin{pmatrix} m_{\text{st}} \sigma_3 & J|\chi| (\cos k_1 + i \cos k_2) \\ J|\chi| (\cos k_1 - i \cos k_2) & -m_{\text{st}} \sigma_3 \end{pmatrix} \begin{pmatrix} f_{e\mathbf{k}} \\ f_{o\mathbf{k}} \end{pmatrix} + \text{const.} \quad (\text{A9})$$

The low-energy Hamiltonian near $(\pm\pi/2, \pi/2)$ points is

$$H \simeq \sum_{\mathbf{k}}' \begin{pmatrix} f_{e1\mathbf{k}}^\dagger & f_{o1\mathbf{k}}^\dagger & f_{e2\mathbf{k}}^\dagger & f_{o2\mathbf{k}}^\dagger \end{pmatrix} \left[\begin{pmatrix} \tau_3 & 0 \\ 0 & -\tau_3 \end{pmatrix} m_{\text{st}}\sigma_3 - J|\chi|k_1 \begin{pmatrix} \tau_1 & 0 \\ 0 & -\tau_1 \end{pmatrix} \right. \\ \left. + J|\chi|k_2 \begin{pmatrix} \tau_2 & 0 \\ 0 & \tau_2 \end{pmatrix} \right] \begin{pmatrix} f_{e1\mathbf{k}} \\ f_{o1\mathbf{k}} \\ f_{e2\mathbf{k}} \\ f_{o2\mathbf{k}} \end{pmatrix}. \quad (\text{A10})$$

Here $f_{e,o1\mathbf{k}}$ describes the field near $(\pi/2, \pi/2)$ and $f_{e,o2\mathbf{k}}$ describes the field near $(-\pi/2, \pi/2)$. The matrices τ_j are Pauli matrices. Note that in this representation $(-\pi/2, -\pi/2)$ and $(\pi/2, -\pi/2)$ are equivalent to $(\pi/2, \pi/2)$ and $(-\pi/2, \pi/2)$, respectively. The Lagrangian is given by

$$L = \int d^2\mathbf{r} \psi^\dagger(\mathbf{r}, t) \left[i\partial_t - J|\chi|i\partial_x \begin{pmatrix} \tau_1 & 0 \\ 0 & -\tau_1 \end{pmatrix} + J|\chi|i\partial_y \begin{pmatrix} \tau_2 & 0 \\ 0 & \tau_2 \end{pmatrix} - m_{\text{st}}\sigma_3 \begin{pmatrix} \tau_3 & 0 \\ 0 & -\tau_3 \end{pmatrix} \right] \psi(\mathbf{r}, t), \quad (\text{A11})$$

where $\psi^\dagger(\mathbf{r}) = (1/\sqrt{\Omega})\sum_k' \exp(-i\mathbf{k} \cdot \mathbf{r}) \begin{pmatrix} f_{e1\mathbf{k}}^\dagger & f_{o1\mathbf{k}}^\dagger & f_{e2\mathbf{k}}^\dagger & f_{o2\mathbf{k}}^\dagger \end{pmatrix}$. After performing the unitary transformation,

$$\psi(\mathbf{r}, t) \rightarrow \begin{pmatrix} 1 & 0 \\ 0 & i\tau_2\sigma_1 \end{pmatrix} \psi(\mathbf{r}, t), \quad (\text{A12})$$

we obtain

$$L = \int d^2\mathbf{r} \bar{\psi}(\mathbf{r}, t) [i\gamma^0\partial_t - J|\chi|\gamma^1i\partial_1 - J|\chi|\gamma^2i\partial_2 - m\sigma_3] \psi(\mathbf{r}, t), \quad (\text{A13})$$

where $\bar{\psi} = \psi^\dagger\gamma^0$,

$$\gamma_0 = \begin{pmatrix} \tau_3 & 0 \\ 0 & -\tau_3 \end{pmatrix}, \quad \gamma_1 = \begin{pmatrix} i\tau_2 & 0 \\ 0 & -i\tau_2 \end{pmatrix}, \quad \gamma_2 = \begin{pmatrix} i\tau_1 & 0 \\ 0 & -i\tau_1 \end{pmatrix}. \quad (\text{A14})$$

Setting $J|\chi| = c_{\text{sw}}$, we obtain

$$S = \int d^3x \bar{\psi}(x) (i\gamma^\mu\partial_\mu - mc_{\text{sw}}\sigma_3) \psi(x), \quad (\text{A15})$$

where $mc_{\text{sw}}^2 = m_{\text{st}}$. Fluctuations about the mean field state are included by a U(1) gauge field, A_μ .^{10,14} This is done by replacing ∂_μ with the covariant derivative $D_\mu = \partial_\mu + iA_\mu$.

The action of the gauge field A_μ is obtained by calculating the fermion polarization function as explicitly shown, for instance, in Ref.¹⁴. Since the fermions are massive, the action has the following Maxwellian form:

$$S_A = -\frac{1}{4e_A^2} \int d^3x F_{\mu\nu} F^{\mu\nu}, \quad (\text{A16})$$

The coefficient $1/e_A^2$ is dependent on the mass through $e_A^2 = 3\pi m$.¹⁴

We derive this action in the imaginary-time formalism. If we take the imaginary time as $it = \tau$, and define $x_3 = c_{\text{sw}}\tau$, then the action takes the so-called Euclidean action:

$$S = \int d^3x \bar{\psi}(x) [\gamma_\mu (\partial_\mu - iA_\mu) + M] \psi(x), \quad (\text{A17})$$

where $M = mc_{\text{sw}}$. The γ -matrices are changed as follows,

$$i\gamma^1 \rightarrow \gamma_1, \quad i\gamma^2 \rightarrow \gamma_2, \quad \gamma^0 \rightarrow \gamma_3. \quad (\text{A18})$$

Their explicit forms are

$$\gamma_1 = \begin{pmatrix} -\tau_2 & 0 \\ 0 & \tau_2 \end{pmatrix}, \quad \gamma_2 = \begin{pmatrix} -\tau_1 & 0 \\ 0 & \tau_1 \end{pmatrix}, \quad \gamma_3 = \begin{pmatrix} \tau_3 & 0 \\ 0 & -\tau_3 \end{pmatrix} \quad (\text{A19})$$

The Dirac fermion Green's function is

$$G_k = -(i\gamma_\mu k_\mu + M)^{-1}. \quad (\text{A20})$$

One obtains,

$$(ik_3\tau_3 - ik_1\tau_2 - ik_2\tau_1 + M)^{-1} = \frac{1}{M^2 + k^2} \begin{pmatrix} -ik_3 + M & k_1 + ik_2 \\ -k_1 + ik_2 & ik_3 + M \end{pmatrix} \quad (\text{A21})$$

Thus, the Green's function is given by

$$G_k = \frac{1}{k^2 + M^2} (ik_\mu \gamma_\mu - M). \quad (\text{A22})$$

Now we integrate out the Dirac fermions fields. After Fourier transformation, the action reads,

$$S = \sum_{k,k'} \overline{\psi_k} [(ik_\mu \gamma_\mu + M) \delta_{k,k'} - i\gamma_\mu A_\mu (k - k')] \psi_{k'}. \quad (\text{A23})$$

Integrating out the Dirac fermions yields

$$\begin{aligned} S &= -Tr \ln (-G_k^{-1}) - Tr \ln [1 + iG\gamma_\mu A_\mu] \\ &= -Tr \ln (-G_k^{-1}) - Tr \left[iG\gamma_\mu A_\mu - \frac{1}{2} (iG\gamma_\mu A_\mu)^2 + \dots \right]. \end{aligned} \quad (\text{A24})$$

The effective action of the gauge field is

$$S^{(2)} = \frac{1}{2} \int \frac{d^3 q}{(2\pi)^3} \Pi_{\mu\nu}(q) A_\mu(-q) A_\nu(q). \quad (\text{A25})$$

The polarization function $\Pi_{\mu\nu}(q)$ is

$$\Pi_{\mu\nu}(q) = - \int \frac{d^3 k}{(2\pi)^3} tr [G_k \gamma_\mu G_{k+q} \gamma_\nu] \quad (\text{A26})$$

Here tr is taken over Dirac fermion components and spin space.

In order to calculate $\Pi_{\mu\nu}(q)$, we use the Feynman trick:

$$\frac{1}{ab} = \int_0^1 dx \frac{1}{[a(1-x) + bx]^2}. \quad (\text{A27})$$

By making use of the Feynman trick, we obtain

$$\begin{aligned} \Pi_{\mu\nu}(q) &= - \int \frac{d^3 k}{(2\pi)^3} \int_0^1 dx \frac{1}{[(k+xq)^2 + x(1-x)q^2 + M^2]^2} \\ &\quad \times tr \left[(ik_\mu \gamma_\mu - M) \gamma_\mu \left(i(k+q)_\mu \gamma_\mu - M \right) \gamma_\nu \right] \\ &= - \int \frac{d^3 k}{(2\pi)^3} \int_0^1 dx \frac{1}{[k^2 + x(1-x)q^2 + M^2]^2} \\ &\quad \times tr \left[\left(i(k-xq)_\mu \gamma_\mu - M \right) \gamma_\mu \left(i(k+(1-x)q)_\mu \gamma_\mu - M \right) \gamma_\nu \right]. \end{aligned} \quad (\text{A28})$$

To calculate the terms including γ -matrices, we use the following formula:

$$\gamma_\mu \gamma_\nu = \delta_{\mu\nu} + i\varepsilon_{\mu\nu\lambda} \tau_\lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (\text{A29})$$

$$tr(\gamma_\mu \gamma_\nu) = \delta_{\mu\nu} tr 1, \quad (\text{A30})$$

$$tr(\gamma_\alpha \gamma_\mu \gamma_\beta \gamma_\nu) = (\delta_{\alpha\mu} \delta_{\beta\nu} - \delta_{\alpha\beta} \delta_{\mu\nu} + \delta_{\alpha\nu} \delta_{\mu\beta}) tr 1. \quad (A31)$$

Thus, we obtain

$$\begin{aligned} \Pi_{\mu\nu}(q) = & -8 \int \frac{d^3 k}{(2\pi)^3} \int_0^1 dx \frac{1}{[k^2 + x(1-x)q^2 + M^2]^2} \\ & \times [-2k_\mu k_\nu + 2x(1-x)(q_\mu q_\nu - q^2 \delta_{\mu\nu}) + [k^2 + x(1-x)q^2 + M^2] \delta_{\mu\nu}], \end{aligned} \quad (A32)$$

where we have used $tr 1 = 8$.

As a convenient calculation scheme, we use the dimensional regularization:³⁹

$$\begin{aligned} \Pi_{\mu\nu}(q) = & -\frac{16\pi^{D/2}}{(2\pi)^D \Gamma(D/2)} \int_0^1 dx \int_0^\infty dk k^{D-1} \frac{2x(1-x)(q_\mu q_\nu - q^2 \delta_{\mu\nu})}{[k^2 + x(1-x)q^2 + M^2]^2} \\ & -8 \int \frac{d^D k}{(2\pi)^D} \int_0^1 dx \frac{[k^2 + x(1-x)q^2 + M^2] \delta_{\mu\nu} - 2k_\mu k_\nu}{[k^2 + x(1-x)q^2 + M^2]^2}. \end{aligned} \quad (A33)$$

The integrations in the first term is calculated as follows

$$\begin{aligned} & \int_0^1 dx x(1-x) \int_0^\infty dk \frac{k^{D-1}}{[k^2 + x(1-x)q^2 + M^2]^2} \\ & = \frac{\pi}{2} \frac{1-D/2}{\sin(\pi D/2)} \int_0^1 dx x(1-x) [M^2 + x(1-x)q^2]^{D/2-2} \end{aligned} \quad (A34)$$

Setting $D = 3$, we obtain

$$\int_0^1 dx x(1-x) [M^2 + x(1-x)q^2]^{-1/2} = \frac{M}{2q^2} + \frac{q^2 - 4M^2}{4q^3} \sin^{-1} \frac{q}{\sqrt{q^2 + 4M^2}}. \quad (A35)$$

The second term is rewritten as

$$\begin{aligned} & \int \frac{d^D k}{(2\pi)^D} \int_0^1 dx \frac{[k^2 + x(1-x)q^2 + M^2] \delta_{\mu\nu} - 2k_\mu k_\nu}{[k^2 + x(1-x)q^2 + M^2]^2} \\ & = \delta_{\mu\nu} \int \frac{d^D k}{(2\pi)^D} \int_0^1 dx \frac{1}{k^2 + x(1-x)q^2 + M^2} - 2 \int \frac{d^D k}{(2\pi)^D} \int_0^1 dx \frac{k_\mu k_\nu}{[k^2 + x(1-x)q^2 + M^2]^2}. \end{aligned} \quad (A36)$$

The integral over k in the second term in the right hand side is carried out as follows

$$\begin{aligned} & \int \frac{d^D k}{(2\pi)^D} \frac{k_\mu k_\nu}{[k^2 + x(1-x)q^2 + M^2]^2} \\ & = \frac{2\pi^{D/2}}{(2\pi)^D D \Gamma(D/2)} \delta_{\mu\nu} \int_0^\infty dk \frac{k^{D+1}}{[k^2 + x(1-x)q^2 + M^2]^2} \\ & = \frac{\pi^{D/2+1}}{2(2\pi)^D \Gamma(D/2)} \frac{[x(1-x)q^2 + M^2]^{D/2-1}}{\sin(\pi D/2)} \delta_{\mu\nu}. \end{aligned} \quad (A37)$$

The first term in the right hand side is,

$$\begin{aligned} & \int \frac{d^D k}{(2\pi)^D} \frac{1}{k^2 + x(1-x)q^2 + M^2} \\ & = \frac{2\pi^{D/2}}{(2\pi)^D \Gamma(D/2)} \int_0^\infty dk \frac{k^{D-1}}{k^2 + x(1-x)q^2 + M^2} \\ & = \frac{\pi^{D/2+1}}{(2\pi)^D \Gamma(D/2)} \frac{[x(1-x)q^2 + M^2]^{D/2-1}}{\sin(\pi D/2)}. \end{aligned} \quad (A38)$$

From these equations, we find that the two terms in the right hand side of Eq. (A36) exactly cancel each other.

As a result, we obtain

$$\Pi_{\mu\nu}(q) = \frac{2}{\pi} (q^2 \delta_{\mu\nu} - q_\mu q_\nu) \left[\frac{M}{2q^2} + \frac{q^2 - 4M^2}{4q^3} \sin^{-1} \frac{q}{\sqrt{q^2 + 4M^2}} \right]. \quad (\text{A39})$$

Expanding the right hand side with respect to q , we obtain

$$\Pi_{\mu\nu}(q) = \frac{1}{3\pi M} (q^2 \delta_{\mu\nu} - q_\mu q_\nu) \left[1 - \frac{1}{10} \left(\frac{q}{M} \right)^2 + \frac{1}{560} \left(\frac{q}{M} \right)^4 + \dots \right]. \quad (\text{A40})$$

On the other hand, the Maxwell term has the following form

$$S_M = \frac{1}{2e_A^2} \sum_q (q^2 \delta_{\mu\nu} - q_\mu q_\nu) A_\mu(-q) A_\nu(q) \quad (\text{A41})$$

Thus, we obtain $e^2 = 3\pi M$.

APPENDIX B: THE BAND NARROWING FACTOR

In this appendix, we evaluate the band narrowing factor omitted in eq.(16). The canonical transformation (9) introduces the following factor in the hopping term,

$$X_i^\dagger X_j = \exp \left[\frac{e_A}{\sqrt{2\Omega}} \sum_q \frac{1}{\omega_q^3} (a_{-q}^\dagger - a_q) e^{iq \cdot R_i} \right] \exp \left[-\frac{e_A}{\sqrt{2\Omega}} \sum_q \frac{1}{\omega_q^3} (a_{-q}^\dagger - a_q) e^{iq \cdot R_j} \right]. \quad (\text{B1})$$

By taking the thermal average with respect to the bosons, we obtain

$$\langle X_i^\dagger X_j \rangle = \exp \left[-\frac{e_A^2}{2\Omega} \sum_q \frac{1}{\omega_q^3} \coth \left(\frac{\beta \omega_q}{2} \right) (1 - \cos[q \cdot (R_i - R_j)]) \right]. \quad (\text{B2})$$

The calculation is similar to eqs.(38) and (39). Thus,

$$\langle X_i^\dagger X_j \rangle \simeq 0.7 \quad (\text{B3})$$

The hopping amplitude is somewhat reduced by the band narrowing factor.

-
- ¹ B. O. Wells, Z.-X. Shen, A. Matsuura, D. M. King, M. A. Kastner, M. Greven, and R. J. Birgeneau, Phys. Rev. Lett. **74**, 964 (1995).
² F. Ronning, C. Kim, D. L. Feng, D. S. Marshall, A. G. Loeser, L. L. Miller, J. N. Eckstein, I. Bozovic, and Z.-X. Shen, Science **282**, 2067 (1998).
³ See, for a review, A. Damascelli, Z. Hussain, and Z. X. Shen, Rev. Mod. Phys. **75**, 473 (2003).
⁴ K.M. Shen, F. Ronning, D.H. Lu, W.S. Lee, N.J.C. Ingle, W. Meevasana, F. Baumberger, A. Damascelli, N.P. Armitage, L.L. Miller, Y. Kohsaka, M. Azuma, M. Takano, H. Takagi, and Z.-X. Shen, Phys. Rev. Lett. **93**, 267002 (2004).
⁵ J. Bała, A. M. Oles, and J. Zaanen, Phys. Rev. B **52**, 4597 (1995).
⁶ T. Xiang and J. M. Wheatley, Phys. Rev. B **54**, R12 653 (1996); B. Kyung and R. A. Ferrell, Phys. Rev. B **54**, 10125 (1996); T. K. Lee and C. T. Shih, Phys. Rev. B **55**, 5983 (1997); T. K. Lee, C.-M. Ho, and N. Nagaosa, Phys. Rev. Lett. **90**, 067001 (2003); T. Tohyama and S. Maekawa, Supercond. Sci. Technol. **13**, R17 (2000).
⁷ Z. Liu and E. Manousakis, Phys. Rev. B **45**, 2425 (1992).
⁸ E. Dagotto, Rev. Mod. Phys. **66**, 763 (1994).
⁹ A. S. Mishchenko and N. Nagaosa, Phys. Rev. Lett. **93**, 036402 (2004).
¹⁰ I. Affleck and J. B. Marston, Phys. Rev. B **37**, R3774 (1988); J. B. Marston and I. Affleck, Phys. Rev. B **39**, 11538 (1989).
¹¹ A. Auerbach and B. E. Larson, Phys. Rev. Lett. **66**, 2262 (1991).
¹² R. B. Laughlin, Phys. Rev. Lett. **79**, 1726 (1997).
¹³ J. B. Marston, Phys. Rev. Lett. **64**, 1166 (1990).

- ¹⁴ D. H. Kim and P. A. Lee, Ann. Phys. **272**, 130 (1999).
- ¹⁵ F. Ronning, C. Kim, Z.-X. Shen, and L. L. Miller, unpublished.
- ¹⁶ M Franz and Z. Tešanović, Phys. Rev. Lett. **87**, 257003 (2001).
- ¹⁷ L. Balents, M. P. A. Fisher, and C. Nayak, Phys. Rev. B **60**, 1654 (1999).
- ¹⁸ Z. Tešanović, O. Vafek and M. Franz, Phys. Rev. B **65**, 180511(R) (2002); I. F. Herbut, Phys. Rev. Lett. **88**, 047006 (2002).
- ¹⁹ I. J. R. Aitchison and N. E. Mavromatos, Phys. Rev. B **53**, 9321 (1996) and references cited therein.
- ²⁰ W. Rantner and X.-G. Wen, Phys. Rev. Lett. **86**, 3871 (2001); Phys. Rev. B **66**, 144501 (2002).
- ²¹ See, also, D. V. Khveshchenko, Phys. Rev. B **65**, 235111 (2002).
- ²² X.-G. Wen and P. A. Lee, Phys. Rev. Lett. **76**, 503 (1996).
- ²³ N. Read and S. Sachdev, Phys. Rev. B **42**, 4568 (1990).
- ²⁴ A. Chubukov, Phys. Rev. B **44**, 12318 (1991).
- ²⁵ C. J. Burden, J. Praschifka, and C. D. Roberts, Phys. Rev. D **46**, 2695 (1992); P. Maris, Phys. Rev. D **52**, 6087 (1995); G. Grignani, G. Semenoff and P. Sodano, Phys. Rev. D **53**, 7157 (1996).
- ²⁶ G. D. Mahan, *Many-Particle Physics* (Plenum, New York, 1990), 2nd edition.
- ²⁷ T. Morinari, cond-mat/0507666, to be published in Phys. Rev. B; cond-mat/0502437.
- ²⁸ J. B. Kogut, Rev. Mod. Phys. **55**, 775 (1983).
- ²⁹ N. Nagaosa and P. A. Lee, Phys. Rev. Lett. **64**, 2450 (1990).
- ³⁰ L. B. Ioffe and P. B. Wiegmann, Phys. Rev. Lett. **65**, 653 (1990).
- ³¹ R. Jackiw and S. Templeton, Phys. Rev. D **23**, 2291 (1981).
- ³² T. W. Appelquist, M. Bowick, D. Karabali and L. C. R. Wijewardhana, Phys. Rev. D **33**, 3704 (1986);
- ³³ T. Appelquist, D. Nash and L. C. R. Wijewardhana, Phys. Rev. Lett. **60**, 2575 (1988).
- ³⁴ K. Kondo and H. Nakatani, Mod. Phys. Lett. **A4**, 2155 (1989).
- ³⁵ B. B. Beard, R. J. Birgeneau, M. Greven, and U. J. Wiese, Phys. Rev. Lett. **80**, 1742 (1998); J.-K. Kim and M. Troyer, Phys. Rev. Lett. **80**, 2705 (1998).
- ³⁶ B. I. Halperin and P. C. Hohenberg, Phys. Rev. **188**, 898 (1969).
- ³⁷ S. Chakravarty, B. I. Halperin, and D. R. Nelson, Phys. Rev. B **39**, 2344 (1989).
- ³⁸ P. Hasenfratz and F. Niedermayer, Z. Phys. B **92**, 91 (1993).
- ³⁹ See, for example, P. Ramond, *Field Theory: A Modern Primer* (Addison-Wesley, 1989).